

ALGEBRAIC PROPERTIES OF QUASI-FINITE COMPLEXES

M. CENCELJ, J. DYDAK, J. SMREKAR, A. VAVPETIČ, AND Ž. VIRK

ABSTRACT. A countable CW complex K is quasi-finite (as defined by A. Karasev [21]) if for every finite subcomplex M of K there is a finite subcomplex $e(M)$ such that any map $f : A \rightarrow M$, where A is closed in a separable metric space X satisfying $X \tau K$, has an extension $g : X \rightarrow e(M)$. Levin's [26] results imply that none of the Eilenberg-MacLane spaces $K(G, 2)$ is quasi-finite if $G \neq 0$. In this paper we discuss quasi-finiteness of all Eilenberg-MacLane spaces. More generally, we deal with CW complexes with finitely many nonzero Postnikov invariants.

Here are the main results of the paper:

Theorem 0.1. *Suppose K is a countable CW complex with finitely many nonzero Postnikov invariants. If $\pi_1(K)$ is a locally finite group and K is quasi-finite, then K is acyclic.*

Theorem 0.2. *Suppose K is a countable non-contractible CW complex with finitely many nonzero Postnikov invariants. If $\pi_1(K)$ is nilpotent and K is quasi-finite, then K is extensionally equivalent to S^1 .*

Date: September 23, 2005.

1991 *Mathematics Subject Classification.* Primary: 54F45; Secondary: 55M10, 54C65 .

Key words and phrases. Extension dimension, cohomological dimension, absolute extensor, universal space, quasi-finite complex, invertible map.

Supported in part by the Slovenian-USA research grant BI-US/05-06/002 and the ARRS research project No. J1-6128-0101-04.

CONTENTS

1. Introduction	2
2. Truncated cohomology	3
3. Homology and cohomology of quasi-finite CW complexes	5
4. Ljubljana complexes	7
5. Application to cohomological dimension theory	9
6. Appendix A	9
7. Appendix B	11
References	12

1. INTRODUCTION

The notation $K \in AE(X)$ or $X\tau K$ means that any map $f : A \rightarrow K$, A closed in X , extends over X .

Theorem 1.1 (Chigogidze). *For each countable simplicial complex P the following conditions are equivalent:*

1. $P \in AE(X)$ implies $P \in AE(\beta(X))$ for any normal space X .
2. There exists a P -invertible map $p : X \rightarrow I^\omega$ of a metrizable compactum X with $P \in AE(X)$ onto the Hilbert cube.

Karasev [21] gave an intrinsic characterization of countable complexes P satisfying 1.1 and called them *quasi-finite complexes*.

Definition 1.2. A CW complex K is called *quasi-finite* if there is a function e from the family of all finite subcomplexes of K to itself satisfying the following property: For every separable metric space X such that K is an absolute extensor of X and for every map $f : A \rightarrow M$, A closed in X , f extends to $g : X \rightarrow e(M)$.

For subsequent generalizations of quasi-finiteness see [22] and [2]. In particular, it is shown in [2] that a countable CW complex K is quasi-finite if and only if $X\tau K$ implies $\beta(X)\tau K$ for all separable metric spaces X . That is an improvement of 1.1.

The first example of a non-quasi-finite CW complex was given by Dranishnikov [9] who showed that $K(Z, 4)$ admits a separable metric space X satisfying $X\tau K(Z, 4)$ but not $\beta(X)\tau K(Z, 4)$ (see [16] for other examples of such X). In [14] it was shown that all $K(G, n)$, $n \geq 3$ and $G \neq 0$, admit a separable metric space X so that $\dim_G(X) = n$ but $\dim_G(\beta(X)) > n$ (see also [23] for related results). Finally, Levin [26] established a result implying the same fact for all $K(G, n)$ so that $G \neq 0$ and $n \geq 2$. The only remaining case among Eilenberg-MacLane spaces are complexes $K(G, 1)$.

Problem 1.3. Characterize groups G such that $K(G, 1)$ is quasi-finite. What are the properties of the class of groups G such that $K(G, 1)$ is quasi-finite?

Problem 1.3 was the main motivation of this paper. More generally, we discuss quasi-finiteness of complexes with finitely many non-trivial Postnikov invariants.

2. TRUNCATED COHOMOLOGY

One of the main tools of this paper is truncated cohomology used for the first time by Dydak and Walsh [17] in their construction of an infinite-dimensional compactum X of integral dimension 2.

Given a pointed CW complex L and a pointed space X we define $h_L^k(X)$ as the $(-k)$ -th homotopy group of the function space $\text{Map}_*(X, L)$, the space of base-point preserving maps whose base-point is the constant map. Since we are interested in Abelian groups, k ranges from minus infinity to -2 . Also, spaces X of interest in this paper are countable CW complexes.

CW complexes L for which truncated cohomology h_L^* is of most use are those with finite homotopy groups. In that case h_L^* is *continuous* in the sense that any map $f : K \rightarrow \Omega^k L$ that is phantom (that means all restrictions $f|_M$ are homotopically trivial for finite subcomplexes M of K) must be homotopically trivial if K is a countable CW complex. In case of L having finite homotopy groups, Levin [25] (see Proposition 2.1) proved that h_L^* is *strongly continuous*: any map $f : N \rightarrow \Omega^k L$, N being a subcomplex of K , that cannot be extended over a countable CW complex K , admits a finite subcomplex M of K such that $f|_{M \cap N}$ cannot be extended over M .

Since we are interested in vanishing of truncated cohomology h_L^* , the remainder of this section is devoted to weak contractibility of mapping spaces.

We first recall a result that in the literature is known as the Zabrodsky Lemma (see Miller [29], Proposition 9.5, and Bousfield [1], Theorem 4.6 as well as Corollary 4.8).

Lemma 2.1. *Let $F \rightarrow E \rightarrow B$ be a fibration where B has the homotopy type of a connected CW complex. Let X be a space. If $\text{Map}_*(F, X)$ is weakly contractible, the induced map $\text{Map}_*(B, X) \rightarrow \text{Map}_*(E, X)$ is a weak homotopy equivalence. ■*

Definition 2.2. Let $\mathcal{P}$ be a set of primes. By a $\mathcal{P}$ -complex we mean a finite CW complex K that is simply connected and all its homotopy groups are $\mathcal{P}$ -groups. That is, homotopy groups of K are finite and the order of each element is a product of primes belonging to $\mathcal{P}$.

A CW complex K is a *co- $\mathcal{P}$ -complex* if for some k the mapping space $\text{Map}_*(\Sigma^k K, L)$ is weakly contractible for all $\mathcal{P}$ -complexes L .

Lemma 2.3. *If K is one of the following*

- (1) *The classifying space BG of a Lie group G with a finite number of path components,*
- (2) *A connected infinite loop space whose fundamental group is a torsion group,*
- (3) *A simply connected CW complex with finitely many homotopy groups,*

then $\text{Map}_(K, L)$ is weakly contractible for all nilpotent finite complexes L with finite homotopy groups.*

Proof. Let L be a finite nilpotent complex with finite homotopy groups. The hypotheses render L complete with respect to Sullivan's finite completion (see [31]). Thus case (1) follows from Friedlander and Mislin [18], Theorem 3.1, while case (2) follows from McGibbon [28], Theorem 3. Case (3) follows from 2.1 and (2) by induction over the number of non-trivial homotopy groups of K . See more details in the proof of 2.8. ■

Proposition 2.4. *A finite product (or a finite wedge) of co- $\mathcal{P}$ -complexes is a co- $\mathcal{P}$ -complex.*

Proof. In case of a finite wedge the proof is quite simple as $\text{Map}_*(K \vee P, L)$ is the product of $\text{Map}_*(K, L)$ and $\text{Map}_*(P, L)$. For the finite product one can use induction plus an observation that 2.1 can be applied to a fibration $F \rightarrow K \rightarrow P$ and yield that K is a co- $\mathcal{P}$ -complex if both F and P are co- $\mathcal{P}$ -complexes. ■

Proposition 2.5. *Let $\mathcal{P}$ be a set of primes. Suppose $K_s, s \in S$, is a family of CW complexes. If there is a natural number k so that all function spaces $\text{Map}_*(\Sigma^k K_s, L)$ are weakly contractible for all $\mathcal{P}$ -complexes L , then the wedge $K = \bigvee_{s \in S} K_s$ is a co- $\mathcal{P}$ -complex. Moreover, if S is countable and each K_s is countable, then the weak product $\prod_{s \in S} K_s$ is a co- $\mathcal{P}$ -complex.*

Proof. The case of the wedge is left to the reader. If S is countable, then each finite product $K_T = \prod_{s \in T} K_s$ has the property that $\text{Map}_*(K_T, \Omega^k L)$ is weakly contractible for any $\mathcal{P}$ -complex L as in the proof of 2.4. Using the fact that truncated cohomology with respect to $\Omega^k L$ is continuous, one gets that $K' = \prod_{s \in S} K_s$, being the direct limit of K_T , also has the property that $\text{Map}_*(K', \Omega^k L)$ is weakly contractible. ■

Definition 2.6. Let $\mathcal{P}$ be a set of primes and let G be a group. G is called a co- $\mathcal{P}$ -group if $K(G, 1)$ is a co- $\mathcal{P}$ -complex.

By Miller's theorem, all locally finite groups are co- $\mathcal{P}$ -groups, where $\mathcal{P}$ is the set of all primes. Another example would consist of all acyclic groups. Divisible groups would serve as well. Note that by the Zabrodsky Lemma 2.1, a group extension $N \rightarrowtail G \rightarrow Q$ implies that under the assumption that N is a co- $\mathcal{P}$ -group, G is a co- $\mathcal{P}$ -group if and only if Q is.

Definition 2.7. Let K be a connected CW complex. We say that K has *finitely many unstable Postnikov invariants* if for some $k \geq 0$, the k -connected cover $K_{<k>}$ of K is an infinite loop space. As usual, $K_{<k>}$ is the (homotopy) fibre of the k -th Postnikov approximation $K \rightarrow P_k(K)$.

Note that infinite loop spaces (in particular infinite symmetric products) and Postnikov pieces are special cases.

Lemma 2.8. *Suppose $\mathcal{P}$ is a set of primes. Let K be a connected CW complex with finitely many unstable Postnikov invariants. K is a co- $\mathcal{P}$ -complex if and only if $G = \pi_1(K)$ is a co- $\mathcal{P}$ -group.*

Proof. Let L be a $\mathcal{P}$ -complex. Let $\tilde{K}$ be the universal cover of K . If K is itself an infinite loop space, so is $\tilde{K}$, and therefore the space $\text{Map}_*(\tilde{K}, L)$ is weakly contractible by Theorem 3 of McGibbon [28]. Otherwise for some $i \geq 1$ the i -connected cover $\tilde{K}_{<i>}$ of $\tilde{K}$ is an infinite loop space. Consider the fibration sequence $\tilde{K}_{<i>} \rightarrow \tilde{K} \rightarrow P_i \tilde{K}$ where $P_i \tilde{K}$ is the i -th Postnikov approximation of $\tilde{K}$. The space $\text{Map}_*(\tilde{K}_{<i>}, L)$ is weakly contractible by Theorem 3 of McGibbon [28]. It follows essentially from Zabrodsky [32], Theorem D, and the fact that L is Sullivan-complete, that the mapping space $\text{Map}_*(P_i \tilde{K}, L)$ is weakly contractible (see also McGibbon [28], Theorem 2). Thus by Lemma 2.1, also the space $\text{Map}_*(\tilde{K}, L)$ is

weakly contractible. The space $\tilde{K}$ sits in the fibration sequence $\tilde{K} \rightarrow K \rightarrow K(G, 1)$ and another application of Lemma 2.1 renders the spaces $\text{Map}_*(K(G, 1), L)$ and $\text{Map}_*(K, L)$ weakly equivalent. $\blacksquare$

Lemma 2.9. *Let $\mathcal{P}$ be a nonempty set of primes. If G is a nilpotent group that is local away from $\mathcal{P}$, then it is a co- $\mathcal{P}$ -group.*

Proof. Let $\mathcal{P}'$ denote the set of primes not in $\mathcal{P}$. The hypotheses on G render $K(G, 1)$ a $\mathcal{P}'$ -local space. By the fundamental theorem of localization of nilpotent spaces it follows that the homology of $K(G, 1)$ is also $\mathcal{P}'$ -local. Let $\cdots \rightarrow L_3 \rightarrow L_2 \rightarrow L_1 \rightarrow L_0$ denote the refined Postnikov tower for L . That is, L_0 is a point and for each i , the fibration $L_i \rightarrow L_{i-1}$ is principal with fibre $K(G_i, k_i)$ where G_i is p -torsion abelian. Note that L is weakly equivalent to the inverse limit $\lim_i L_i$, and since $K(G, 1)$ is a CW complex it suffices to show that $\text{Map}_*(K(G, 1), \lim_i L_i)$ is weakly contractible. This latter space is homeomorphic with the inverse limit $\lim_i \text{Map}_*(K(G, 1), L_i)$. Since the fibrations are principal, the Puppe sequence shows that we only need to consider reduced cohomology $\tilde{H}^*(K(G, 1); G_i)$ with coefficients in G_i . Since $H_*(G)$ is local away from $\mathcal{P}$ it follows by the universal coefficient theorem that $\tilde{H}^*(K(G, 1); G_i)$ is trivial. $\blacksquare$

Corollary 2.10. *Suppose $\mathcal{P}$ is a set of primes and G is a nilpotent group with Abelianization $\text{Ab}(G)$. If $\text{Ab}(G)/\text{Tor}(\text{Ab}(G))$ is $\mathcal{P}$ -divisible, then G is a co- $\mathcal{P}$ -group.*

Proof. By 6.4 $\text{Ab}(G)/\text{Tor}(\text{Ab}(G))$ is $\mathcal{P}$ -divisible if and only if G is local away from $\mathcal{P}$. $\blacksquare$

3. HOMOLOGY AND COHOMOLOGY OF QUASI-FINITE CW COMPLEXES

In this section we deal with (co)homological properties of quasi-finite complexes. First, we need a generalization of Theorem II of [15].

Theorem 3.1. *Suppose K is a countable CW complex and h_* is a generalized reduced homology theory such that $h_*(K) = 0$. For any CW complex P and any $\alpha \in h_*(P) \setminus \{0\}$ there is a compactum X and a map $f : A \rightarrow P$ from a closed subset A of X such that $X \tau K$, $\alpha = f_*(\gamma)$ for some $\gamma \in \check{h}_*(A)$ and γ is 0 in $\check{h}_*(X)$.*

Proof. Replacing P by the carrier of α we may assume P is finite. Compactum X is built as in Theorem II of [15]. We start with $X_1 = \text{Cone}(P)$, $A_1 = P$ and build an inverse sequence (X_n, A_n) of compact polyhedra so that for every extension problem $g : B \rightarrow K$, B closed in X_n , there is $m > n$ and a map $G : X_m \rightarrow K$ extending $g \circ p_n^m : B' \rightarrow K$, where $p_n^m : X_m \rightarrow X_n$ is the bonding map and $B' = (p_n^m)^{-1}(B)$. For each n we have $\gamma_n \in h_*(A_n)$ which vanishes in $h_*(X_n)$. In the inductive step we pick an extension problem $g : B \rightarrow K$, B closed in X_n , create an extension $G : X_n \rightarrow \text{Cone}(K)$, and consider the pull-back E of the projection $K \times I \rightarrow \text{Cone}(K)$ under G . The projection $p : E \rightarrow X_n$ has fibers being either homeomorphic to K or single points. Therefore $h_*(p)$ is an isomorphism and one can pick a finite subpolyhedron A_{n+1} of E carrying $\gamma_{n+1} \in h_*(A_{n+1})$ which gets mapped to γ_n under $h_*(p)$. Since γ_{n+1} vanishes in $h_*(E)$, it vanishes in a finite subpolyhedron X_{n+1} of E containing A_{n+1} . Since there are only countably many extension problems to be solved (see [8] or [10]) that process produces an inverse sequence whose inverse limit (X, A) satisfies $X \tau K$ and one has $\gamma \in \check{h}_*(A)$ so that γ vanishes in $\check{h}_*(X)$ and $f_*(\gamma) = \alpha$, where $f : A \rightarrow P = A_1$ is the projection. $\blacksquare$

Theorem 3.2. *Suppose K is a countable CW complex and h^* is a strongly continuous truncated cohomology theory such that $h^*(K) = 0$. For any countable CW complex P and any $\alpha \in h^*(P) \setminus \{0\}$ there is a compactum X and a map $f : A \rightarrow P$ from a closed subset A of X such that $X\tau K$ and there is no $\gamma \in \check{h}^*(X)$ satisfying $\gamma|_A = f^*(\alpha)$.*

Proof. We can reduce the proof to the case of P being a finite polyhedron as there is a finite subcomplex M of P so that $\alpha|_M \neq 0$ and that M can be used instead of P . Compactum X is built as in 3.1. We start with $X_1 = \text{Cone}(P)$, $A_1 = P$ and built an inverse sequence (X_n, A_n) of compact polyhedra so that for every extension problem $g : B \rightarrow K$, B closed in X_n , there is $m > n$ and a map $G : X_m \rightarrow K$ extending $g \circ p_n^m : B' \rightarrow K$, where $p_n^m : X_m \rightarrow X_n$ is the bonding map and $B' = (p_n^m)^{-1}(B)$. Also, for each n the pullback α_n of α under $A_n \rightarrow A_1$ does not extend over X_n . In the inductive step we pick an extension problem $g : B \rightarrow K$, B closed in X_n , create an extension $G : X_n \rightarrow \text{Cone}(K)$, and consider the pull-back E of the projection $K \times I \rightarrow \text{Cone}(K)$ under G . The projection $p : E \rightarrow X_n$ has fibers being either homeomorphic to K or single points. Therefore $p^* = h^*(p)$ is an isomorphism. Since $p^*(\alpha_n)$ does not extend over E , there is a finite subpolyhedron X_{n+1} of E such that $p^*(\alpha_n)$ restricted to $A_{n+1} = X_{n+1} \cap p^{-1}(A_n)$ does not extend over X_{n+1} . Since there are only countably many extension problems to be solved (see [8] or [10]) that process produces an inverse sequence whose inverse limit (X, A) satisfies $X\tau K$ and the projection $f : A \rightarrow P = A_1$ has the property that there is no $\gamma \in \check{h}^*(X)$ satisfying $\gamma|_A = f^*(\alpha)$. ■

Recall that, given a map $i : M \rightarrow N$, $X\tau i$ means that for any map $f : A \rightarrow M$, A closed in X , there is a map $g : X \rightarrow N$ extending $i \circ f$.

Theorem 3.3. *Suppose K is a countable CW complex and $i : M \rightarrow N$ is a map of CW complexes such that $X\tau K$ implies $X\tau i$ for all compacta X .*

- (1) *If h_* is a generalized reduced homology theory such that the inclusion induced homomorphism $h_*(M) \rightarrow h_*(N)$ is not trivial, then $h_*(K) \neq 0$.*
- (2) *If h^* is a truncated strongly continuous cohomology theory such that the inclusion induced homomorphism $h^*(N) \rightarrow h^*(M)$ is not trivial and M is countable, then $h^*(K) \neq 0$.*

Proof. We may assume i is an inclusion.

1. Suppose $\alpha \in h_*(M)$ does not become 0 in $h_*(N)$. As in 3.1 pick a map $f : A \rightarrow M$ of a closed subset of a compactum X so that $X\tau K$ and γ equals 0 in $\check{h}_*(e(M))$ for some $\gamma \in \check{h}_*(A)$ satisfying $f_*(\gamma) = \alpha$. If f extends to $g : X \rightarrow N$, then $\alpha = f_*(\gamma)$ becomes 0 in $h_*(N)$, a contradiction.

2. Suppose $\alpha \in h^*(N)$ and $\alpha|_M \neq 0$. We may reduce this case to M finite by switching to a finite subcomplex L of M with the property $\alpha|_L \neq 0$. As in 3.2 pick a map $f : A \rightarrow M$ of a closed subset of a compactum X so that $X\tau K$ and $f^*(\alpha|_M)$ does not extend over X . If $f : A \rightarrow M$ extends to $g : X \rightarrow N$, then $g^*(\alpha) \in \check{h}^*(X)$ extends $f^*(\alpha|_M)$, a contradiction. ■

Theorem 3.4. *Suppose $\mathcal{P}$ is a set of primes. Let K be a connected countable co- $\mathcal{P}$ -complex. If K is quasi-finite, then it is $Z_{(\mathcal{P})}$ -acyclic.*

Proof. Assume K is quasi-finite and not $Z_{(\mathcal{P})}$ -acyclic. Replace K with ΣK (using [2]) if necessary to ensure $H_k(K; Z_{(\mathcal{P})}) \neq 0$ for some $k \geq 2$. Let $\alpha_K \in H_k(K; Z_{(\mathcal{P})})$ be nonzero. Since K is the colimit of its finite subcomplexes, α_K is

the image of $\alpha_M \in H_k(M; Z_{(\mathcal{P})})$ for some finite subcomplex M of K . Certainly the image of α_M under $H_k(M; Z_{(\mathcal{P})}) \rightarrow H_k(e(M); Z_{(\mathcal{P})})$ is nontrivial. Thus Lemma 7.2 yields a $\mathcal{P}$ -complex L with the restriction morphism $[e(M), \Omega^2 L] \rightarrow [M, \Omega^2 L]$ nontrivial. This is to say that $h^*(e(M)) \rightarrow h^*(M)$ is nontrivial where h^* is the truncated cohomology theory defined by virtue of $\Omega^2 L$. The hypotheses on L ensure strong continuity of h^* . Thus the nontriviality of $h^*(e(M)) \rightarrow h^*(M)$ contradicts (2) of Theorem 3.3. ■

Corollary 3.5. *Let K be a countable CW complex with finitely many nonzero homotopy groups and $G = \pi_1(K)$ nilpotent. Suppose that G is not torsion. If K is quasi-finite, the group $FG = G/\text{Tor}(G)$ (and thus also $\text{Ab}(G)/\text{Tor}(\text{Ab}(G))$) is not divisible by any prime p .*

Proof. Suppose that, on the contrary, FG is divisible by a prime p , hence local away from p . Since G is not torsion and is nilpotent, also $\text{Ab}(G)$ is not torsion, hence certainly $H_1(K) \otimes Z_{(p)}$ is nontrivial. Thus Theorem 3.4 yields a contradiction. ■

3.4 and 2.8 imply the following.

Corollary 3.6. *Let K be a simply connected countable CW complex with at least one and at most finitely many nontrivial homotopy groups. Then K is not quasi-finite.* ■

Corollary 3.7. *Suppose G is a locally finite countable group. If $K(G, 1)$ is quasi-finite, then G is acyclic.* ■

However, there are some countable acyclic groups G for which $K(G, 1)$ are also not quasi-finite. Cencelj and Repovš [3], using results of Dranishnikov and Repovš [13] showed in §5 that the minimal grope M^* which is $K(\pi_1(M^*), 1)$ is not quasi-finite. This holds also for the fundamental group of any grope: For a grope M let $\gamma(m)$ denote the maximal number of handles on the discs with handles used in the construction of the m -stage of M . Modify the inverse limit construction of the example of [3] replacing every simplex in the triangulation of the k -th element of the inverse system by the n -th stage of the grope which has every generator replaced by a disc with $\gamma(kn)$ handles.

4. LJUBLJANA COMPLEXES

Definition 4.1. A connected CW complex L is called a *Ljubljana complex* if there is a co- $\mathcal{AP}$ -complex K , $\mathcal{AP}$ being the set of all primes, such that, for any compactum X , the conditions $X\tau L$ and $X\tau K(H_1(K), 1)$ imply $X\tau K$.

Lemma 4.2. *Suppose $F \rightarrow E \rightarrow B$ is a fibration of connected CW complexes. If F is a co- $\mathcal{AP}$ -complex, $\mathcal{AP}$ being the set of all primes, and B is a Ljubljana complex, then E is a Ljubljana complex.*

Proof. Notice $\pi_1(E) \rightarrow \pi_1(B)$ is an epimorphism (use the long exact sequence of a fibration) which implies $H_1(E) \rightarrow H_1(B)$ is an epimorphism.

Pick a co- $\mathcal{AP}$ -complex K such that $X\tau K$ and $X\tau K(H_1(B), 1)$ imply $X\tau B$ for all compacta X . Let M be the wedge of F , K , $K(Q, 1)$, and $X\tau K(Z/p^\infty, 1)$ for all primes p . By 2.5 and the Miller Theorem, M is a co- $\mathcal{AP}$ -complex. Suppose X is a compactum such that $X\tau M$ and $X\tau K(H_1(E), 1)$. By 6.5 one gets $X\tau K(H_1(B), 1)$ which, together with $X\tau K$, implies $X\tau B$. Since $X\tau F$ and $X\tau B$, we infer $X\tau E$. ■

Corollary 4.3. *Let L be a connected CW complex with nilpotent fundamental group. If L has finitely many unstable Postnikov invariants, then L is a Ljubljana complex.*

Proof. Notice that the universal cover $\tilde{L}$ of L is a co- $\mathcal{AP}$ -complex by 2.8. We get $K(\pi_1(L), 1)$ is a Ljubljana complex by 6.6. The fibration $\tilde{L} \rightarrow L \rightarrow K(\pi_1(L), 1)$ implies L is a Ljubljana complex. ■

Definition 4.4. A connected CW complex L is called *extensionally Abelian* if $X\tau K(H_n(L), n)$ for all $n \geq 1$ imply $X\tau K$ for all compacta X .

Proposition 4.5. *Each extensionally Abelian complex L is a Ljubljana complex.*

Proof. Let K be the weak product of $K(H_n(L), n)$, $n \geq 2$. By (2) of 2.3, K is a co- $\mathcal{AP}$ -complex. Clearly, $X\tau K$ and $X\tau K(H_1(L), 1)$ imply $X\tau K(H_n(L), n)$ for all $n \geq 1$. Thus $X\tau L$. ■

Proposition 4.6. *A finite wedge (or finite product) of Ljubljana complexes is a Ljubljana complex.*

Proof. Let L be the wedge (or the product) of Ljubljana complexes L_s , $s \in S$, where S is finite. For each $s \in S$ choose a co- $\mathcal{AP}$ -complex K_s such that for any compactum X the conditions $X\tau K_s$ and $X\tau K(H_1(L_s), 1)$ imply $X\tau L_s$. Let K be the wedge of all K_s . By 2.4 it is a co- $\mathcal{AP}$ -complex. Notice that $H_1(L_s)$ is a retract of $H_1(L)$ for each $s \in S$. Therefore any compactum X satisfying

- a. $X\tau K(H_1(L), 1)$,
- b. $X\tau K$,

also satisfies $X\tau K(H_1(L_s), 1)$ for each $s \in S$. Hence $X\tau L_s$ for each $s \in S$ which implies $X\tau L$. ■

There is a connection between Ljubljana complexes and co- $\mathcal{P}$ -complexes.

Proposition 4.7. *Suppose K is a countable Ljubljana complex. If $\mathcal{P}$ is a set of primes such that $H_1(K)/\text{Tor}(H_1(K))$ is $\mathcal{P}$ -divisible, then K is a co- $\mathcal{P}$ -complex.*

Proof. Choose a co- $\mathcal{AP}$ -complex L such that, for any compactum X , the conditions $X\tau L$ and $X\tau K(H_1(K), 1)$ imply $X\tau K$. Let $\mathcal{P}'$ be the complement of $\mathcal{P}$ in the set of all primes. Consider K' , the wedge of L , $K(Z_{(\mathcal{P}')} , 1)$, $K(Q, 1)$, and all $K(Z/p, 1)$ (p ranging through all primes). By 2.10 and 2.5 K' is a co- $\mathcal{P}$ -complex. Since $X\tau K'$ implies $X\tau K$ for all compacta, 3.3 implies that there is $k \geq 0$ such that the truncated cohomology of K with respect to $\Omega^k L$, L any $\mathcal{P}$ -complex, is trivial. Thus K is a co- $\mathcal{P}$ -complex. ■

Theorem 4.8. *Suppose K is a countable Ljubljana complex such that $\Sigma^m K$ is equivalent (over the class of compacta) to a quasi-finite countable complex L for some $m \geq 0$. If K is not acyclic, then it is equivalent to S^1 .*

Proof. We may assume L is simply connected as $\Sigma^{m+1} K$ is equivalent to ΣL (see [11]) and ΣL is quasi-finite by [2].

Suppose K is not equivalent to S^1 . Choose a co- $\mathcal{AP}$ -complex P such that conditions $X\tau P$ and $X\tau K(H_1(K), 1)$ imply $X\tau K$. Let $k \geq 2$ be a number such that all maps $\Sigma^n P \rightarrow R$ are null-homotopic if R is an $\mathcal{AP}$ -complex and $n \geq k$.

Step 1. L is not contractible as otherwise $\Sigma^m K$ would have to be contractible implying K being acyclic.

Step 2. L is not acyclic as it is not contractible by Step 1.

Step 3. Since $X\tau K$ implies $X\tau K(H_1(K), 1)$, the group $H_1(K)$ has the property of $H_1(K)/\text{Tor}(H_1(K))$ being divisible by some prime p . Indeed, If $H_1(K)/\text{Tor}(H_1(K))$ is not divisible by any prime, then the Bockstein basis of $H_1(K)$ consists of all Bockstein groups and $X\tau K(H_1(K), 1)$ implies $X\tau S^1$ by Bockstein First Theorem. Since $X\tau K$ implies $X\tau K(H_1(K), 1)$ and $X\tau S^1$ implies $X\tau K$ for any compactum X , K is equivalent to S^1 over compacta.

Let e be the function of L .

Case 1: $H_*(K)$ is a torsion group. There is M such that $H_*(M) \rightarrow H_*(e(M))$ is not trivial. By 7.1, there is a map $f : \Sigma^k(e(M)) \rightarrow J$ such that $f|_{\Sigma^k(M)}$ is not trivial, J is simply connected, and all homotopy groups of J are finite. Consider the wedge N of P and $K(\bigoplus_q Z/q, 1)$. Notice $X\tau N$ implies $X\tau K(H_1(K), 1)$. Therefore $X\tau N$ implies $X\tau K$ which, in turn, implies $X\tau L$ and $X\tau i_M$, where $i_M : M \rightarrow e(M)$. Since $\text{Map}_*(N, \Omega^k(J))$ is weakly contractible, 3.3 implies homotopy triviality of $f|_{\Sigma^k(M)}$, a contradiction.

Case 2: $H_*(K)$ is not a torsion group. Notice $H_*(L)$ is not torsion as well. Indeed if $H_*(L)$ is torsion, we could find a finite dimensional compactum Y of high rational dimension but all torsion dimensions equal 1. Such compactum satisfies $Y\tau L$ but $Y\tau \Sigma^m K$ fails as it implies rational dimension of Y to be at most $m + n$, where $H_n(K)$ is not torsion. There is M such that the image of $H_*(M) \rightarrow H_*(e(M))$ is not torsion. Therefore there is $n > 0$ such that $H_n(M; Z_{(p)}) \rightarrow H_n(e(M); Z_{(p)})$ is not trivial. By 7.2, there is a map $f : \Sigma^k(e(M)) \rightarrow J$ such that $f|_{\Sigma^k(M)}$ is not trivial, J is simply connected, and all homotopy groups of J are finite p -groups. Consider the wedge N of P and $K(Z[\frac{1}{p}] \oplus Z/p, 1)$. Using 6.7 and 6.5, one gets $X\tau N$ implies $X\tau K$ which, in turn, implies $X\tau L$ and $X\tau i_M$, where $i_M : M \rightarrow e(M)$. Since $\text{Map}_*(N, \Omega^k(J))$ is weakly contractible, 3.3 implies homotopy triviality of $f|_{\Sigma^k(M)}$, a contradiction. ■

Corollary 4.9. *Suppose G is a nontrivial nilpotent group. If $K(G, 1)$ is quasi-finite, then it is equivalent, over the class of paracompact spaces, to S^1 .*

5. APPLICATION TO COHOMOLOGICAL DIMENSION THEORY

Theorem 5.1. *Suppose $G \neq 1$ is a countable group such that $\dim_G(\beta(X)) = 1$ for every separable metric space X satisfying $\dim_G(X) = 1$. If G is nilpotent, then $\dim_G(X) \leq 1$ implies $\dim(X) \leq 1$ for all paracompact spaces X .*

Proof. By an improvement of Chigogidze's Theorem 1.1 contained in [2], $K(G, 1)$ is quasi-finite. Therefore 4.9 says $K(G, 1)$ is equivalent to S^1 over compacta. A result in [2] says that $K(G, 1)$ is equivalent to S^1 over paracompact spaces which completes the proof. ■

6. APPENDIX A

In this section we discuss results related to groups that are needed in the paper.

Lemma 6.1. *Let p be a natural number and let $\mathcal{D}_p$ be the class of groups G such that $\text{Ab}(G)/\text{Tor}(\text{Ab}(G))$ is p -divisible, where $\text{Ab}(G)$ is the Abelianization of G . If $f : G \rightarrow H$ is an epimorphism and $G \in \mathcal{D}_p$, then $H \in \mathcal{D}_p$.*

Proof. Notice that $G \in \mathcal{D}_p$ if and only if for each $a \in G$ there is $b \in G$ and $k \geq 1$ such that $(a \cdot b^{-p})^k$ belongs to the commutator subgroup $[G, G]$ of G . Suppose $a \in H$. Pick $b \in G$ with $a = f(b)$. There is $c \in G$ and $k \geq 1$ such that $(b \cdot c^{-p})^k \in [G, G]$. Now $(a \cdot f(c)^{-p})^k \in [H, H]$ and $H \in \mathcal{D}_p$. ■

Lemma 6.2. *Let p be a natural number and let $\mathcal{D}_p$ be the class of groups G such that $H/\text{Tor}(H)$ is p -divisible, where H is the Abelianization of G . If G, H are Abelian and $G \in \mathcal{D}_p$, then $G \otimes H \in \mathcal{D}_p$.*

Proof. It suffices to show that for each element a of $G \otimes H$ there is $b \in G \otimes H$ and an integer $k \neq 0$ such that $k \cdot a + kp \cdot b = 0$. That in turn can be reduced to generators of $G \otimes H$ of the form $g \otimes h$. Pick $u \in G$ and an integer $k \neq 0$ such that $k \cdot g + kp \cdot u = 0$. Now $k \cdot (g \otimes h) + kp \cdot (u \otimes h) = 0$. ■

We recall a result of Robinson (see [30], 5.2.6) on the relation between a nilpotent group and its abelianization.

Proposition 6.3 (Robinson). *Let $\mathcal{N}$ denote the category of nilpotent groups. Let $\mathcal{P}$ be a class of groups in $\mathcal{N}$ with the following properties.*

- (1) *For A and B abelian, $B \in \mathcal{P}$, any quotient of $A \otimes B$ belongs to $\mathcal{P}$.*
- (2) *For $K, Q \in \mathcal{P}$, an extension $1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1$ in $\mathcal{N}$ implies $G \in \mathcal{P}$.*

Suppose that $G \in \mathcal{N}$. If $\text{Ab}(G)$ belongs to $\mathcal{P}$, so does G . ■

We note the following corollary.

Corollary 6.4. *Let G be a nilpotent group and set $H = \text{Ab}(G)$. If $H/\text{Tor}(H)$ is p -divisible, so is $G/\text{Tor}(G)$.*

Proof. Define the class $\mathcal{D}_p$ by letting a nilpotent group G belong to $\mathcal{D}_p$ if and only if $F_p(G) = G/\text{Tor}_p(G)$ is p -divisible where $\text{Tor}_p(G)$ denotes the p -torsion subgroup of G . Note that $F_p(G)$ is p -divisible if and only if $G/\text{Tor}(G)$ is, hence it suffices to check properties (1) and (2) of Proposition 6.3.

As for (1) it follows from 6.1 and 6.2.

For (2), note that F_p is a functor $\mathcal{N} \rightarrow \mathcal{N}$. Let $1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1$ be an extension in $\mathcal{N}$. We apply F_p . Since $\text{Tor}_p(K) = K \cap \text{Tor}_p(G)$, the morphism $F_p(K) \rightarrow F_p(G)$ is injective. Evidently, $q: F_p(G) \rightarrow F_p(Q)$ is surjective. Moreover, $F_p(K)$ is a subset of the kernel of q . Assume that K belongs to $\mathcal{D}_p$. If $q(\xi) = 1$ for some $\xi \in F_p(G)$, then $\xi^{p^i} \in F_p(K)$ for large enough i . By assumption on K , the group $F_p(K)$ is p -divisible, hence $\xi^{p^i} = \eta^{p^i}$ for an element $\eta \in F_p(K)$. But $F_p(G)$ is free of p -torsion (and nilpotent), so the equality $\xi^{p^i} = \eta^{p^i}$ in $F_p(G)$ implies $\xi = \eta$ (see for example Hilton, Mislin, Roitberg [20], Corollary 2.3). Therefore in fact $\xi \in F_p(K)$, ie $\ker q = F_p(K)$. This is to say that $1 \rightarrow F_p(K) \rightarrow F_p(G) \rightarrow F_p(Q) \rightarrow 1$ is an extension in $\mathcal{N}$. If, in addition, Q belongs to $\mathcal{D}_p$ then $F_p(K)$ and $F_p(Q)$ are p -divisible and free of p -torsion, and as such local away from p . Therefore so is $F_p(G)$, by Corollary 2.5 of [20]. ■

Lemma 6.5. *Let $f: G \rightarrow H$ be an epimorphism of Abelian groups and let X be a compactum. If*

- a. $X\tau K(G, 1)$,
- b. $X\tau K(Q, 1)$,
- c. $X\tau K(Z/p^\infty, 1)$ for all primes p ,

then $X\tau K(H, 1)$.

Proof. Suppose $X\tau K(H, 1)$ fails. This can only happen if there is a Bockstein group F in the Bockstein basis $\sigma(H)$ such that $\dim_F(X) > 1$. That group must be either $Z_{(p)}$ or Z/p for some p . $Z_{(p)}$ belongs to $\sigma(H)$ if and only if $H/\text{Tor}(H)$ is not divisible by p in which case $Z_{(p)}$ belongs to $\sigma(G)$ by 6.1 and $\dim_F(X) \leq 1$ by Bockstein First Theorem. Therefore $F = Z/p$ which means that $\text{Tor}(H)$ is not divisible by p . Now, either G is not divisible by p or its torsion group is not divisible by p implying $\dim_F(X) \leq 1$, a contradiction. ■

Corollary 6.6. *Let G be a nilpotent group with Abelianization $\text{Ab}(G)$ and let X be a compactum. If*

- a. $X\tau K(\text{Ab}(G), 1)$,
- b. $X\tau K(Q, 1)$,
- c. $X\tau K(Z/p^\infty, 1)$ for all primes p ,

then $X\tau K(G, 1)$.

Proof. Consider the lower central series of G : $G = \Gamma^1 G \supset \Gamma^2 G \supset \dots \supset \Gamma^i G \supset \dots$. Let $F_i = \Gamma^i G / \Gamma^{i+1} G$. Since there is an epimorphism from $F_i \otimes \text{Ab}(G)$ to F_{i+1} , where $\text{Ab}(G)$ is the Abelianization of G , $X\tau K(F_i, 1)$ for all i by 6.5. We proceed by induction on $c - i$ (c being the nilpotency class of G) showing that $X\tau K(\Gamma^i G, 1)$. If $c - i$ is 0, then $\Gamma^i G = F_i$ and we are done. Since the sequence $1 \rightarrow \Gamma^{i+1} G \rightarrow \Gamma^i G \rightarrow F_i \rightarrow 1$ is exact, one uses a fibration $K(\Gamma^{i+1} G, 1) \rightarrow K(\Gamma^i G, 1) \rightarrow K(F_i, 1)$ to conclude $X\tau K(\Gamma^i G, 1)$ given $X\tau K(\Gamma^{i+1} G, 1)$. That constitutes the inductive step and, as $\Gamma^1 G = G$, we get $X\tau K(G, 1)$. ■

Corollary 6.7. *Let p be a natural number and let $\mathcal{D}_p$ be the class of groups G such that $H/\text{Tor}(H)$ is p -divisible, where H is the Abelianization of G . If $G \in \mathcal{D}_p$ is nilpotent and $X\tau K(Z[\frac{1}{p}] \oplus Z/p, 1)$, then $X\tau K(G, 1)$ for any compactum X .*

Corollary 6.8. *Let G be a nilpotent group. If the Abelianization $\text{Ab}(G)$ of G is a torsion group and $X\tau K(\bigoplus_p Z/p, 1)$, then $X\tau K(G, 1)$ for any compactum X .*

7. APPENDIX B

In this Appendix we prove results allowing us to detect homology via maps to finite complexes with finite homotopy groups.

Lemma 7.1. *Let A be a finite CW complex and $\alpha \in H_k(A)$ a nontrivial element where $k \geq 2$. There exists a finite $(k - 1)$ -connected CW complex B with finite homotopy groups and a map $f: A \rightarrow B$ with $\beta = f_*(\alpha)$ nontrivial. Furthermore, if α is of infinite order in $H_k(A)$, then β may be assumed to be of order r for any given natural $r \geq 2$.*

Proof. With the exception of the statements about the connectedness and the order, this is precisely Lemma 2.1 of Levin [26]. In the course of proving the cited lemma, Levin constructs a $(k - 1)$ -connected complex L , and he makes β of order 2 if α has infinite order. The generalization to arbitrary r is trivial. ■

Lemma 7.2. *Let M be a finite CW complex, and let $\mathcal{P}$ be a nonempty set of primes. Let $\alpha \in H_k(M; Z_{(\mathcal{P})})$ be a nontrivial element for some $k \geq 2$. Then there exists a finite $(k - 1)$ -connected CW complex N with $\mathcal{P}$ -torsion homotopy groups and a map $f: M \rightarrow N$ with $f_*(\alpha)$ nontrivial.*

Proof. The assumption is that there exists an element $\alpha \in H_k(M)$ which is either $\mathcal{P}$ -torsion or it has infinite order. We can apply Lemma 7.1 to obtain a $(k-1)$ -connected finite complex with finite homotopy groups N' and a map $f': M \rightarrow N'$ with $\beta' = f'_*(\alpha)$ nontrivial of order all of whose prime divisors belong to $\mathcal{P}$. Let $N' \rightarrow N$ be localization at the set $\mathcal{P}$. Then β' will map to nontrivial β under localization $\tilde{H}_*(N') \rightarrow \tilde{H}_*(N) = \tilde{H}_*(N') \otimes Z_{(\mathcal{P})}$ and N is (homotopy equivalent to) the finite complex as in the statement of the lemma. ■

REFERENCES

- [1] A.K.Bousfield, *Localization and periodicity in unstable homotopy theory*, J. Amer. Math. Soc. **7** (1994), no. 4, 831–873.
- [2] M. Cencelj, J. Dydak, J. Smrekar, A. Vavpetič, and Ž. Virk, *Compact maps and quasi-finite complexes*, preprint.
- [3] M.Cencelj and D. Repovš, *On compacta of cohomological dimension one over nonabelian groups*, Houston J. Math. **26**(2000), 527–536.
- [4] A.Chigogidze, *Compactifications and universal spaces in extension theory*, arXiv:math.GN/9908073 v1 15 Aug 1999.
- [5] A.Chigogidze, *Cohomological dimension of Tychonov spaces*, Topology Appl. **79** (1997), 197–228.
- [6] A.Chigogidze, *Infinite dimensional topology and Shape theory*, in: Handbook of Geometric Topology, ed. by R. Daverman and R. Sher.
- [7] A.Chigogidze, *Inverse Spectra*, North Holland, Amsterdam, 1996.
- [8] A.Dranishnikov, *On extension theory of compact spaces*,
- [9] A.N.Dranishnikov, *Cohomological dimension is not preserved by Stone- Čech compactification*, Comptes Rendus Bulgarian Acad. Sci. **41** (1988), 9–10.
- [10] A.N.Dranishnikov, *Cohomological Dimension Theory of Compact Metric Spaces*, Topology Atlas (1999).
- [11] A. N. Dranishnikov and J. Dydak, *Extension dimension and extension types*, Proc. Steklov Math. Inst. **212** (1996), 55–88.
- [12] A.Dranishnikov and J.Dydak, *Extension theory of separable metrizable spaces with applications to dimension theory*, Transactions of the American Math.Soc. **353** (2000), 133–156.
- [13] A.Dranishnikov and D.Repovš, *Cohomological dimension with respect to perfect groups*, Topology and its Applications **74** (1996), 123–140.
- [14] J.Dydak, *Compactifications and cohomological dimension*, Topology and its Appl. **50** (1993), 1–10.
- [15] J.Dydak, *Realizing dimension functions via homology*, Topology and its Appl. **64** (1995), 1–7.
- [16] J.Dydak, J.J.Walsh, *Spaces without cohomological dimension preserving compactifications*, Proc. Amer. Math. Soc. **113** (1991), 1155–1162.
- [17] J.Dydak, J.J.Walsh, *Infinite dimensional compacta having cohomological dimension two: An application of the Sullivan Conjecture*, Topology **32** (1993), 93–104.
- [18] E. M. Friedlander, G. Mislin, *Locally finite approximation of Lie groups. I*, Invent. Math. **83** (1986), no. 3, 425–436.
- [19] A.Hatcher, *Algebraic Topology*, Cambridge University Press, Cambridge, 2002.
- [20] P. Hilton, G. Mislin, J. Roitberg, *Localization of Nilpotent groups and spaces*, North-Holland Publishing Co. , Amsterdam-Oxford; American Elsevier Publishing Co. , Inc. , New York, 1975.
- [21] A.Karasev, *On two problems in extension theory*, arXiv:math.GT/0312269 v1 12 Dec 2003.
- [22] A.Karasev and V.Valov, *Extension dimension and quasi-finite CW complexes*, preprint, 2004.
- [23] A.I.Karinski, *On cohomological dimension of the Stone- Čech compactification*, Vestnik Moscow Univ. no. 4 (1991), 8–11 (in Russian).
- [24] W.I.Kuzminov, *Homological dimension theory*, Russian Math. Surveys **23** (1968), 1–45.
- [25] M.Levin, *Constructing Compacta of Different Extensional Dimensions*, Canad. Math. Bull. Vol. **44** (1) (2001), pp. 80–86.
- [26] M.Levin, *Some examples in cohomological dimension theory*, Pacific Journal of Mathematics **202** (2002), 371–378.
- [27] S.Mardešić and J.Segal, *Shape theory*, North-Holland Publ.Co. (1982).

- [28] C.A.McGibbon, *A note on Miller's theorem about maps out of classifying spaces*, Proc. Amer. Math. Soc. 124 (1996), no. 10, 3241–3245.
- [29] H.Miller, *The Sullivan conjecture on maps from classifying spaces*, Ann. of Math. (2) 120(1) (1984), 39–87.
- [30] D. J. S. Robinson, *A course in the theory of groups*, Springer-Verlag, New York, 1993.
- [31] D.Sullivan, *Genetics of homotopy theory and the Adams conjecture*, Ann. of Math. (2) 100 (1974), 1–79.
- [32] A.Zabrodsky, *On phantom maps and a theorem of H. Miller*, Israel J. Math 58 (1987), no. 2, 129–143.

FAKULTETA ZA MATEMATIKO IN FIZIKO, UNIVERZA V LJUBLJANI, JADRANSKA ULICA 19, SI-1111 LJUBLJANA, SLOVENIJA

E-mail address: `matija.cencelj@guest.arnes.si`

UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996, USA

E-mail address: `dydak@math.utk.edu`

DEPARTAMENT D'ÀLGEBRA I GEOMETRIA, UNIVERSITAT DE BARCELONA, GRAN VIA DE LES CORTS CATALANES, 585, E-08007 BARCELONA, ESPAÑA

E-mail address: `smrekar@ub.edu`

FAKULTETA ZA MATEMATIKO IN FIZIKO, UNIVERZA V LJUBLJANI, JADRANSKA ULICA 19, SI-1111 LJUBLJANA, SLOVENIJA

E-mail address: `ales.vavpetic@fmf.uni-lj.si`

FAKULTETA ZA MATEMATIKO IN FIZIKO, UNIVERZA V LJUBLJANI, JADRANSKA ULICA 19, SI-1111 LJUBLJANA, SLOVENIJA

E-mail address: `ziga.virk@student.fmf.uni-lj.si`